Opiate use statistics

- 669,000 Americans reported using heroin in the past year.
- Twenty-six (26) % of persons seeking treatment for SA/SD reported opiates as the primary substance (16% heroin, 10% prescription).
- Persons meeting the DSM Criteria has doubled (100%) since 2007.
- Over half of all young persons receiving treatment for heroin dependence started by abusing prescription opioids (replicated in 3 recent independent surveys).
- Unintentional poisoning from prescription opioids quadrupled from 1999-2010
- This death rate now outpaces the rates of death from both heroin and cocaine combined.

<u>Profile of Today's Opioid Dependent Person</u> (Treatment Episode Date Set 2002-2012, DHHS)

- White, non-Hispanic (65% of heroin admissions); 86% for opiate, non-heroin
 - ✓ Black persons btw the ages of 20 to 34 accounted for 1/10 of 1% of admissions.
- Male (41% vs. 24% for females).
- Young Adults
 - ✓ Forty-seven percent of admissions for opiates other than heroin were aged 20 to 29 (2012).
 - ✓ Fifty-five percent of admissions for heroin were age 20 to 34 (2012).
- Peak age for all opiate treatment seekers is 25 years of age.
- Using pills in late teens.
 - ✓ First use of prescription opiates is predominantly after the age of 16 (79%) compared to 50% for other substances.
- Has likely had at least five (5) treatment episodes.
- Injecting (71%).

Opiate Treatment Principles

- Addiction is a Brain Disease
- Opiate Addiction is highly influenced by endorphin deprivation
- Structured Program of Recovery: Every day of abstinence improves the person's likelihood of long term brain healing and personal recovery.
- Post Acute Withdrawal:
 - ✓ can last up to 18 months
 - ✓ is a major reason for relapse (unstable brain chemistry leading to unstable thought, actions and behaviors)
 - ✓ Although not a psychiatric disorder, PAWS has symptoms that mimic mental illness.

Treatment Phases for Opiate Dependence

Transition Phase
Initiation Phase
Implementation Phase
Stabilization Phase Half Way House Option
Stabilization Phase Outpatient Option

Stabilization Phase Half Way House and Outpatient Option Maintenance Phase